



6-2016

## Some Results on f-Simultaneous Chebyshev Approximation

Sh. Al-Sharif  
*Yarmouk University*

Kh. Qaraman  
*Yarmouk University*

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Numerical Analysis and Computation Commons](#)

---

### Recommended Citation

Al-Sharif, Sh. and Qaraman, Kh. (2016). Some Results on f-Simultaneous Chebyshev Approximation, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 11, Iss. 1, Article 22.  
Available at: <https://digitalcommons.pvamu.edu/aam/vol11/iss1/22>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact [hvkoshy@pvamu.edu](mailto:hvkoshy@pvamu.edu).



## Some Results on $f$ -Simultaneous Chebyshev Approximation

Sh. Al-Sharif and Kh. Qaraman

Mathematics Department  
Yarmouk University  
Irbid, Jordan

[sharifa@yu.edu.jo](mailto:sharifa@yu.edu.jo), [math811@yahoo.com](mailto:math811@yahoo.com)

Received: February 26, 2015; Accepted: October 30, 2015

### Abstract

Let  $X$  be a Hausdorff topological vector space and  $f$  be a real valued continuous function on  $X$ . In this paper we introduce and study the concept of  $f$ -simultaneous approximation of a nonempty subset  $K$  of  $X$  as a generalization to the problem of simultaneous approximation. Further we present some results regarding  $f$ -simultaneous approximation in the quotient space.

**Keywords:** Hausdorff topological vector space;  $f$ -best simultaneous approximation;  $f$ -simultaneous Chebyshev; simultaneous approximation; quotient space

**MSC 2010 No.:** 41A65, 41A50

### 1. Introduction

Let  $K$  be a subset of a Hausdorff topological vector space  $X$  and  $f$  be a real valued continuous function on  $X$ . For  $x \in X$ , set  $F_K(x) = \inf_{k \in K} f(x - k)$ . A point  $k_0 \in K$  is called  $f$ -best approximation to  $x$  in  $K$  if  $F_K(x) = f(x - k_0)$ . The set  $P_K^f(x) = \{k_0 \in K : F_K(x) = f(x - k_0)\}$  denotes the set of all  $f$ -best approximations to  $x$  in  $K$ . Note that this set may be empty. The set  $K$  is said to be  $f$ -proximal ( $f$ -Chebyshev) if for each  $x \in X$ ,  $P_K^f(x)$  is non-empty (singleton). The notion of  $f$ -best approximation in a vector space  $X$  was given by Breckner and Brosowski and in a Hausdorff topological vector space  $X$  by Narang. For a Hausdorff locally convex topological vector space and a continuous sublinear functional  $f$  on  $X$ ,

Breckner, Brosowski, and Govindarajulu proved certain results on best approximation relative to the functional  $f$ . By using the existence of elements of  $f$ -best approximation some results on fixed point were proved by Pai and Veermani.

As a generalization to the problem of simultaneous approximation (see Saidi and Singer), we introduce the concept of best  $f$ -simultaneous approximation as follows:

**Definition 1.**

Let  $f$  be a real valued continuous function on a Hausdorff topological vector space  $X$ . A subset  $A$  of  $X$  is called  $f$ -bounded if there exists  $M > 0$  such that  $|f(x)| \leq M$  every  $x \in A$ .

Note that  $f$ -bounded sets need not be bounded in the classical sense, for example if  $f(x) = e^{-x}$ , the set  $[0, \infty)$  is an  $f$ -bounded subset of real numbers.

**Definition 2.**

Let  $X$  be a Hausdorff topological real vector space,  $f$  be a real valued continuous function on  $X$ , and  $K$  be a non-empty subset of  $X$ . A point  $k_0 \in K$  is called  $f$ -best simultaneous approximation in  $K$  if there exists an  $f$ -bounded subset  $A$  of  $X$  such that

$$F_K(A) = \inf_{k \in K} \sup_{a \in A} |f(a - k)| = \sup_{a \in A} |f(a - k_0)|.$$

The set of all  $f$ -best simultaneous approximations to an  $f$ -bounded subset  $A$  of  $X$  in  $K$  is denoted by

$$P_K^f(A) = \left\{ k \in K : F_K(A) = \sup_{a \in A} |f(a - k)| \right\}.$$

The set  $K$  is called  $f$ -simultaneously proximal ( $f$ -simultaneously Chebyshev) if for each  $f$ -bounded set  $A$  in  $X$ ,  $P_K^f(A) \neq \emptyset$  (singleton).

We note that if  $f(x) = \|x\|$  ( $f(x) = \|x\| + \epsilon$ ), then the concept of  $f$ -best approximation is precisely best approximation, i.e. best  $\epsilon$ -approximation (see Khalil, Rezapour, Singer and others).

A set  $K$  is said to be inf-compact at a point  $x \in X$ , (see Pai and Veermani), if each minimizing sequence in  $K$  (i.e.  $f(x - k_n) \rightarrow F_K(x)$ ) has a convergent subsequence in  $K$ . The set  $K$  is called inf-compact if it is inf-compact at each  $x \in X$ . A subset  $K$  of  $X$  is called  $f$ -compact, (see Moghaddam), if for every sequence  $\{k_n\}$  in  $K$ , there exist a subsequence  $\{k_{n_i}\}$  of  $\{k_n\}$  and  $k_0 \in K$  such that  $f(k_{n_i} - k_0) \rightarrow 0$ . It is easy to see that if  $K$  is  $f$ -compact or inf-compact, then  $K$  is  $f$ -simultaneously proximal.

In this paper we introduce and study the concept of  $f$ -simultaneous approximation of a subspace  $K$  of a Hausdorff topological real vector space  $X$ , and existence and uniqueness. Certain results regarding  $f$ -simultaneous approximation in quotient spaces is obtained by generalizing some of the results in Moghaddam.

Throughout this paper  $X$  is a Hausdorff topological real vector space and  $f$  is a real valued continuous function on  $X$ .

## 2. $f$ -Simultaneous Approximation

In this section we give some characterization of  $f$ -proximal sets in  $X$ . We begin with the following definitions:

### Definition 3.

A function  $f : X \rightarrow \mathbb{R}$  is called

- (1) absolutely subadditive if  $|f(x+y)| \leq |f(x)| + |f(y)|$  for all  $x, y \in X$ .
- (2) absolutely homogeneous if  $f(\alpha x) = |\alpha| f(x)$ , for all  $x \in X$  and all  $\alpha \in \mathbb{R}$ .

### Definition 4.

A subset  $K$  of  $X$  is called  $f$ -closed if for all sequences  $\{k_m\}$  of  $K$  and for all  $x \in X$  such that  $f(x - k_m) \rightarrow 0$ , we have  $x \in K$ .

### Theorem 1.

Let  $K$  be a subset of  $X$ . Then,

- (1)  $F_{K+y}(A+y) = F_K(A)$ , for all  $f$ -bounded sets  $A \subset X$ ,  $y \in X$ .
- (2)  $P_{K+y}^f(A+y) = P_K^f(A) + y$ , for all  $f$ -bounded sets  $A \subset X$ ,  $y \in X$ .
- (3)  $K$  is  $f$ -simultaneously proximal ( $f$ -simultaneously Chebyshev) if and only if  $K+y$  is  $f$ -simultaneously proximal ( $f$ -simultaneously Chebyshev) for every  $y \in X$ .

Moreover if  $f$  is an absolutely homogeneous function, then

- (4)  $F_{\lambda K}(\lambda A) = |\lambda| F_K(A)$ , for all  $f$ -bounded sets  $A \subset X$  and  $\lambda \in \mathbb{R}$ .
- (5)  $P_{\lambda K}^f(\lambda A) = \lambda P_K^f(A)$ , for all  $f$ -bounded sets  $A \subset X$  and  $\lambda \in \mathbb{R}$ .
- (6)  $K$  is  $f$ -simultaneously proximal ( $f$ -simultaneously Chebyshev) if and only if  $\lambda K$  is  $f$ -simultaneously proximal ( $f$ -simultaneously Chebyshev),  $\lambda \in \mathbb{R}$ .

### Proof:

- (1) Let  $A \subset X$ ,  $f$ -bounded set. Then

$$F_{K+y}(A+y) = \inf_{w \in K} \sup_{a \in A} |f((a+y) - (w+y))| = F_K(A).$$

- (2) The equation

$$\sup_{a \in A} |f(a - k_0)| = \inf_{k \in K} \sup_{a \in A} |f((a+y) - (k+y))| = \inf_{k \in K} \sup_{a \in A} |f(a - k)|,$$

implies that  $k_0 + y \in P_{K+y}^f(A+y)$  if and only if  $k_0 \in P_K^f(A)$ . Thus

$$P_{K+y}^f(A+y) = P_K^f(A) + y.$$

- (3) This follows immediately from part two.

- (4) Let  $A \subset X$  be an  $f$ -bounded set,  $\lambda \in \mathbb{R}$ . Then

$$F_{\lambda K}(\lambda A) = \inf_{k \in K} \sup_{a \in A} |f(\lambda a - \lambda k)| = |\lambda| \inf_{k \in K} \sup_{a \in A} |f(a - k)| = |\lambda| F_K(A).$$

(5) If  $\lambda = 0$ , we are done. If  $\lambda \neq 0$  and  $k_0 \in P_{\lambda K}^f(\lambda A)$ , then  $k_0 \in \lambda K$  and

$$\sup_{a \in A} |f(\lambda a - k_0)| = \inf_{k \in K} \sup_{a \in A} |f(\lambda a - \lambda k)|.$$

This implies that

$$\sup_{a \in A} \left| f\left(a - \frac{1}{\lambda} k_0\right) \right| = F_K(A),$$

which implies that  $\frac{1}{\lambda} k_0 \in P_K^f(A)$ .

(6) This follows immediately from part 5.  $\square$

## Theorem 2.

Let  $f$  be an absolutely homogeneous real valued function on  $X$  and  $M$  be a subspace of  $X$ . Then,

(1)  $F_M(\lambda A) = |\lambda| F_M(A)$ , for all  $f$ -bounded sets  $A \subset X$  and  $\lambda \in \mathbb{R} - \{0\}$ .

(2)  $P_M^f(\lambda A) = \lambda P_M^f(A)$ , for all  $f$ -bounded sets  $A \subset X$  and  $\lambda \in \mathbb{R} - \{0\}$ .

**Proof:**

(1) Let  $A \subset X$  be an  $f$ -bounded set and  $\lambda \neq 0 \in \mathbb{R}$ . Then,

$$F_M(\lambda A) = \inf_{m \in M} \sup_{a \in A} |f(\lambda a - m)| = |\lambda| \inf_{m' \in M} \sup_{a \in A} |f(a - m')| = |\lambda| F_M(A).$$

(2) Let  $m_0 \in P_M^f(\lambda A)$ . Then,

$$\begin{aligned} \sup_{a \in A} |\lambda| \left| f\left(a - \frac{1}{\lambda} m_0\right) \right| &= \sup_{a \in A} |f(\lambda a - m_0)| \\ &= \inf_{m \in M} \sup_{a \in A} |f(\lambda a - m)| \\ &= \inf_{m' \in M} \sup_{a \in A} |\lambda| \left| f(a - m') \right|. \end{aligned}$$

Therefore,

$$\sup_{a \in A} \left| f\left(a - \frac{1}{\lambda} m_0\right) \right| = \inf_{m' \in M} \sup_{a \in A} |f(a - m')| = F_M(A),$$

for all  $\lambda \in \mathbb{R} - \{0\}$ , which implies that  $\frac{1}{\lambda} m_0 \in P_M^f(A)$ , and so  $m_0 \in \lambda P_M^f(A)$ .  $\square$

For a subset  $K$  of  $X$ , let us define  $\widehat{K}_F$  such that

$$\widehat{K}_F = \left\{ A \subset X : F_K(A) = \sup_{a \in A} f(a) \right\}.$$

Using this we prove the following theorem characterizing  $f$ -simultaneously proximal sets.

## Theorem 3.

Let  $K$  be a subspace of  $X$ . Then  $K$  is  $f$ -simultaneously proximal in  $X$  if and only if every  $f$ -bounded subset  $A$  of  $X$  can be written as  $B + k$  for some  $k \in K$  and  $B \in \widehat{K}_F$ .

**Proof:**

Suppose the condition hold. Let  $A \subset X$  be an  $f$ -bounded subset of  $X$ . By assumption there exists  $k_0 \in K$  and  $B \in \widehat{K}_F$  such that  $A = B + k_0$ . Hence  $A - k_0 \in \widehat{K}_F$ . Therefore,

$$\begin{aligned} \sup_{a \in A} |f(a - k_0)| &= F_K(A - k_0) \\ &= \inf_{k \in K} \sup_{a \in A} |f(a - k_0 - k)| \\ &= \inf_{k' \in K} \sup_{a \in A} |f(a - k')| = F_K(A). \end{aligned}$$

Hence,  $K$  is  $f$ -simultaneously proximal.

Conversely, suppose  $K$  is  $f$ -simultaneously proximal and  $A \subset X$  be an  $f$ -bounded subset of  $X$ . Then there exists  $k_0 \in K$  such that

$$\sup_{a \in A} |f(a - k_0)| = \inf_{k \in K} \sup_{a \in A} |f(a - k)| = \inf_{k' \in K} \sup_{a \in A} |f(a - (k' + k_0))|$$

where  $k = k' + k_0$ . Hence,

$$\sup_{a \in A} |f(a - k_0)| = F_K(A - k_0).$$

Consequently,  $A - k_0 \in \widehat{K}_F$ . So there exists  $B \in \widehat{K}_F$  such that  $A - k_0 = B$  or  $A = B + k_0$ .  $\square$

**Theorem 4.**

Let  $f$  be a real valued continuous function on  $X$  such that  $x = 0$  if and only if  $f(x) = 0$ . If  $K$  is  $f$ -simultaneously proximal, then  $K$  is  $f$ -closed.

**Proof:**

Let  $\{k_m\}$  be a sequence of  $K$  and  $x \in X$ , such that  $f(x - k_m) \rightarrow 0$ . Taking  $A = \{x\}$ , we have

$$F_K(A) = \inf_{k \in K} \sup_{a \in A} |f(a - k)| \leq |f(x - k_m)| \rightarrow 0.$$

Since  $K$  is  $f$ -simultaneously proximal, there exists  $k_0 \in K$  such that

$$F_K(A) = |f(x - k_0)| = 0.$$

Hence,  $f(x - k_0) = 0$ . Using assumption it follows that  $x - k_0 = 0$ . Therefore,  $x = k_0 \in K$  and  $K$  is  $f$ -closed.  $\square$

**3.  $f$ -Simultaneous Approximation in Quotient Space**

Let  $M$  be a closed subspace of  $X$ . Then a function  $\tilde{f}: (X/M) \rightarrow \mathbb{R}$  can be defined as follows:

$$\tilde{f}(x + M) = \inf_{y \in M} |f(x + y)|.$$

**Proposition 1.**

Let  $M$  be a closed subspace of  $X$ . If  $A$  is  $f$ -bounded in  $X$ , then  $A/M$  is  $\tilde{f}$ -bounded in  $X/M$ .

**Proof:**

Let  $A$  be an  $f$ -bounded subset in  $X$ . Since  $M$  is a subspace, for  $x + M \in A/M$

$$\left| \tilde{f}(x + M) \right| = \inf_{y \in M} |f(x + y)| \leq |f(x)|.$$

Consequently since  $A$  is an  $f$ -bounded subset of  $X$ , it follows that  $A/M$  is  $\tilde{f}$ -bounded in  $X/M$ .  $\square$

**Theorem 5.**

Let  $M$  a closed subspace of  $X$ . If  $B$  is  $\tilde{f}$ -bounded in  $X/M$ , then there exists an  $f$ -bounded subset  $A$  of  $X$  such that  $B = A/M$ .

**Proof:**

Let  $B$  be a nonempty  $\tilde{f}$ -bounded in  $X/M$ . Let  $C = \bigcup_{b \in B} b$ .

Claim:  $B = \{\bar{x} = x + M : x \in C\}$ . Indeed if  $b \in B$ , then  $b = x_b + M$  for some  $x_b \in X$ . But  $M$  is a subspace. Thus  $x_b = x_b + 0 \in x_b + M \subset C$ . Hence  $b = x_b + M \in \{\bar{x} = x + M : x \in C\}$  and  $B \subseteq \{\bar{x} = x + M : x \in C\}$ . Similarly if  $x \in C$ , then  $x \in b_x + M$  for some  $b_x + M \in B$ . This implies that  $x = b_x + m_x$  for some  $m_x \in M$ . Hence  $x + M = b_x + m_x + M = b_x + M \in B$ . Therefore,  $\{\bar{x} = x + F : x \in C\} \subseteq B$ .

Now clearly  $C$  is not bounded unless  $M$  is trivial. Note that  $B$  is  $\tilde{f}$ -bounded. So there exists  $K > 0$  such that  $\left| \tilde{f}(b) \right| \leq K$  for all  $b \in B$ . Consider the set  $A = \{x \in C : |f(x)| \leq K + 1\} \subseteq C$ .

Now we claim that for all  $x \in C$ ,

$$\bar{x} \cap A = (x + M) \cap A \neq \phi.$$

Given  $x \in C$ . Since

$$\left| \tilde{f}(x + M) \right| = \inf_{m \in M} |f(x + m)| \leq K,$$

there exists  $m_x \in M$  such that  $|f(x + m_x)| < K + 1$ . But  $x + m_x \in x + M \subseteq C$ . Hence  $x + m_x \in (x + M) \cap A \neq \phi$ . Claim  $B = A/M$ . Since  $A \subseteq C$ , we have  $A/M \subseteq \{\bar{x} = x + M : x \in C\} = B$ . To show the other inclusion, let  $b \in B = \{\bar{x} = x + M : x \in C\}$ . Then  $b = x_b + M$  for some  $x_b \in C$ . But  $(x_b + M) \cap A \neq \phi$ . Thus there exists  $a \in A$  such that  $a = x_b + m_a \in x_b + M$ . Therefore,  $b = x_b + M = (x_b + m_a) + M = a + M \in A/M$ . Hence  $B \subseteq A/M$ . Consequently  $A/M = B$ .  $\square$

**Theorem 6.**

Let  $K$  be a subspace of  $X$  and  $M$  be a closed  $f$ -proximal subspace of  $K$ . If  $k_0$  is a point of  $f$ -best simultaneous approximation to  $A \subset X$  in  $K$ , then  $k_0 + M$  is an  $\tilde{f}$ -best simultaneous approximation to  $A/M$  in  $K/M$ .

**Proof:**

Suppose  $k_0 + M$  is not  $\tilde{f}$ -best simultaneous approximation to  $A/M$  in  $K/M$ . Then, for at least  $k \in K$ , say  $k_1 \in K$ , we have

$$\sup_{a \in A} \tilde{f}(a - k_1 + M) < \sup_{a \in A} \tilde{f}(a - k_0 + M).$$

Since

$$\sup_{a \in A} \tilde{f}(a - k_0 + M) = \sup_{a \in A} \inf_{m \in M} |f(a - k_0 + m)| \leq \sup_{a \in A} |f(a - k_0)|,$$

we have

$$\sup_{a \in A} \tilde{f}(a - k_1 + M) = \sup_{a \in A} \inf_{m \in M} |f(a - k_1 + m)| < \sup_{a \in A} |f(a - k_0)|.$$

But  $M$  is  $f$ -proximal, so for some  $m_0 \in M$  we have

$$\sup_{a \in A} |f(a - k_1 + m_0)| = \sup_{a \in A} \inf_{m \in M} |f(a - k_1 + m)| < \sup_{a \in A} |f(a - k_0)|.$$

Since  $M \subset K$ , it follows that  $k_1 - m_0 \in K$ . Therefore,  $k_0$  not  $f$ -best simultaneous approximation to  $A$  in  $K$ , which is a contradiction.  $\square$

**Corollary 1.**

Let  $K$  be a subspace of  $X$  and  $M$  is a closed  $f$ -proximal subspace of  $K$ . If  $K$  is  $f$ -simultaneously proximal in  $X$ , then  $K/M$  is  $\tilde{f}$ -simultaneously proximal in  $X/M$ .

**Proof:**

Let  $B$  be an  $\tilde{f}$ -bounded subset of  $X/M$ . Then, by Theorem 5, there exists  $f$ -bounded subset  $A \subset X$  such that  $B = A/M$ . If  $K$  is  $f$ -simultaneously proximal in  $X$ , then there exists at least  $k_0 \in K$  such that  $k_0$  is  $f$ -best simultaneous approximation to  $A$  in  $K$ . By Theorem 6,  $k_0 + M$  is an  $\tilde{f}$ -best simultaneous approximation to  $A/M$  in  $K/M$ , so  $K/M$  is  $\tilde{f}$ -simultaneously proximal in  $X/M$ .  $\square$

**Theorem 7.**

Let  $K$  be a subspace of  $X$  and  $M$  is a closed  $f$ -proximal subspace of  $K$ . If  $K/M$  is  $\tilde{f}$ -simultaneously proximal in  $X/M$ , then  $K$  is  $f$ -simultaneously proximal in  $X$ .

**Proof:**

Let  $A$  be an  $f$ -bounded subset of  $X$ . By Proposition 1,  $A/M$  is  $\tilde{f}$ -bounded in  $X/M$ . Since  $K/M$  is  $\tilde{f}$ -simultaneously proximal in  $X/M$ , then there exists  $k_0 + M \in K/M$  such that  $k_0 + M$  is  $\tilde{f}$ -best simultaneous approximation to  $A/M$  from  $K/M$ , so

$$\begin{aligned} \sup_{a \in A} \tilde{f}(a - k_0 + M) &= \inf_{k \in K} \sup_{a \in A} \tilde{f}(a - k + M) \\ &= \inf_{k \in K} \sup_{a \in A} \inf_{m \in M} |f(a - k + m)| \\ &\leq \inf_{k \in K} \sup_{a \in A} |f(a - k + m)| \\ &= \inf_{k \in K} \sup_{a \in A} |f(a - k')|, \end{aligned} \tag{1}$$



where,  $k' = k - m \in K$ . Since  $M$  is  $f$ -proximal, there exists  $m_0 \in M$  such that

$$\sup_{a \in A} |f(a - k_0 - m_0)| = \sup_{a \in A} \inf_{m \in M} |f(a - k_0 + m)| = \sup_{a \in A} \tilde{f}(a - k_0 + M). \quad (2)$$

Consequently, combining (1) and (2) since  $M \subset K$ , it follows that

$$\begin{aligned} \sup_{a \in A} |f(a - k_0 - m_0)| &\leq \inf_{k' \in K} \sup_{a \in A} |f(a - k')| \\ &\leq \sup_{a \in A} |f(a - k_0 - m_0)| \end{aligned}$$

Hence,

$$\sup_{a \in A} |f(a - k_0 + m_0)| = \inf_{k \in K} \sup_{a \in A} |f(a - k)|$$

So  $k_0 + m_0$  is an  $f$ -best simultaneous approximation to  $A$  from  $K$  and  $K$  is  $f$ -simultaneously proximal in  $X$ .  $\square$

### Theorem 8.

Let  $W$  and  $M$  be two subspaces of  $X$ . If  $M$  is a closed  $f$ -proximal subspace of  $X$ , then the following assertions are equivalent:

- (1)  $W/M$  is  $\tilde{f}$ -simultaneously proximal in  $X/M$ ,
- (2)  $W + M$  is  $f$ -simultaneously proximal in  $X$ .

**Proof:**

(1)  $\Rightarrow$  (2). Since  $(W+M)/M = W/M$  and  $M$  are  $f$ -simultaneously proximal, using Theorem 7, it follows that  $W + M$  is  $f$ -simultaneously proximal in  $X$ .

(2)  $\Rightarrow$  (1). Since  $W + M$  is  $f$ -simultaneously proximal and  $M \subseteq W + M$ , by Corollary 1,  $(W + M)/M = W/M$  is simultaneously  $\tilde{f}$ -proximal.  $\square$

### Theorem 9.

Let  $K, M$  be two subspaces of  $X$  such that,  $M \subset K$ . If  $M$  is closed  $\tilde{f}$ -simultaneously proximal in  $X$  and  $K$  is  $f$ -simultaneously Chebyshev in  $X$ , then,  $K/M$  is  $\tilde{f}$ -simultaneously Chebyshev in  $X/M$ .

**Proof:**

Suppose not. Then there exists  $A$ ,  $f$ -bounded subset of  $X$  such that  $A/M \in X/M$  is  $\tilde{f}$ -bounded and  $k_1 + M, k_2 + M \in P_{K/M}^{\tilde{f}}(A/M)$  such that  $k_1 + M \neq k_2 + M$ . Thus  $k_1 - k_2 \notin M$ . Since  $M$  is an  $f$ -simultaneously proximal in  $X$ , then

$$P_M^f(A - k_1) \neq \phi, \text{ and } P_M^f(A - k_2) \neq \phi.$$

Let  $m_1 \in P_M^f(A - k_1)$ , and  $m_2 \in P_M^f(A - k_2)$ . By Theorem 7,  $k_1 + m_1$  and  $k_2 + m_2$  are  $f$ -best simultaneous approximations to  $A$  from  $K$ . Since  $K$  is  $f$ -simultaneously Chebyshev in  $X$ , then  $k_1 + m_1 = k_2 + m_2$  and hence  $k_1 - k_2 = m_1 - m_2 \in M$ , which is a contradiction.  $\square$

**Definition 5.**

A subset  $K$  of  $X$  is called  $f$ -quasi-simultaneously Chebyshev if  $P_K^f(A)$  is nonempty and  $f$ -compact set in  $X$  for all  $f$ -bounded subsets of  $X$ .

**Theorem 10.**

Let  $M$  be a closed  $f$ -simultaneously proximal subspace of  $X$  and  $K$  is  $f$ -quasi-simultaneously Chebyshev of  $X$  such that  $M \subset K$ . Then,  $K/M$  is  $\tilde{f}$ -quasi-simultaneously Chebyshev in  $X/M$ .

**Proof:**

Since  $K$  is  $f$ -simultaneously proximal in  $X$ , By Corollary 1,  $K/M$  is  $\tilde{f}$ -simultaneously proximal in  $X/M$ . Let  $B$  be an  $\tilde{f}$ -bounded subset of  $X/M$ . Then, by Theorem 5,  $B = A/M$  for an  $f$ -bounded subset  $A$  of  $X$ . If  $(k_n + M)$  a sequence in  $P_{K/M}^{\tilde{f}}(A/M)$ , by the proof of Theorem 7, for every  $n$ , there exists  $m_n \in M$  such that  $k_n + m_n = k'_n \in P_K^f(A)$ . But since  $M$  is a subspace, we have

$$k'_n + M = k_n + m_n + M = k_n + M.$$

Since  $K$  is  $f$ -quasi-simultaneously Chebyshev in  $X$ , the sequence  $\{k_n\}$  has a subsequence  $\{k_{n_i}\}$  such that  $f(k_{n_i} - k_0) \rightarrow 0$  for some  $k_0 \in P_K^f(A)$ . But

$$\tilde{f}(k_{n_i} - k_0 + M) \leq |f(k_{n_i} - k_0)| \rightarrow 0.$$

Therefore,

$$\tilde{f}(k_{n_i} - k_0 + M) \rightarrow 0$$

and

$$\tilde{f}((k_{n_i} + M) - (k_0 + M)) \rightarrow 0.$$

Hence,  $P_{K/M}^{\tilde{f}}(A/M)$  is  $\tilde{f}$ -compact and  $K/M$  is  $\tilde{f}$ -quasi-simultaneously Chebyshev. This complete the proof.  $\square$

**Definition 6.**

A topological vector space  $X$  is said to have the  $f$ -property if every  $f$ -bounded sequence in  $X$  has an  $f$ -convergent subsequence, where  $f$  is a real valued continuous function on  $X$ .

Note that the space  $X = l^2$  has the  $f$ -property for every projection  $f : X \rightarrow \mathbb{R}$ , and if  $f(x) = \|x\|$ , then every finite dimensional Banach space has the  $f$ -property.

**Proposition 2.**

Let  $f$  be an absolutely homogeneous subadditive continuous real valued function on a topological vector space  $X$  and  $K$  be an  $f$ -closed subspace of  $X$ . Then, for any  $f$ -bounded subset  $A$  of  $X$ ,  $P_K^f(A)$  is  $f$ -closed.

**Proof:**

Let  $K$  be an  $f$ -closed subspace of  $X$  and  $A$  be an  $f$ -bounded subset of  $X$ . If  $\{k_m\}$  is a sequence in  $P_K^f(A)$  and  $x \in X$  such that  $f(k_m - x) \rightarrow 0$ , then  $x \in K$  since  $K$  is  $f$ -closed.

Further,

$$\begin{aligned}\inf_{k \in K} \sup_{a \in A} |f(a - k)| &= \sup_{a \in A} |f(a - k_m)| \\ &= \sup_{a \in A} |f((a - x) - (k_m - x))| \\ &\geq \sup_{a \in A} ||f(a - x)| - |f(k_m - x)||.\end{aligned}$$

Taking the limit as  $m \rightarrow \infty$ , we get

$$\inf_{k \in K} \sup_{a \in A} |f(a - k)| \geq \sup_{a \in A} |f(a - x)|.$$

Consequently,

$$\inf_{k \in K} \sup_{a \in A} |f(a - k)| = \sup_{a \in A} |f(a - x)|.$$

Then,  $x \in P_K^f(A)$  and  $P_K^f(A)$  is  $f$ -closed.  $\square$

### Theorem 11.

Let  $f$  be a real valued sub-additive continuous function on a topological vector space  $X$  that has the  $f$ -property and  $M$  be a closed subspace of  $X$ . If  $W$  is a subspace of  $X$  such that  $W + M$  is  $f$ -closed, then the following assertions are equivalent:

- (1)  $W/M$  is  $\tilde{f}$ -simultaneously quasi-Chebyshev in  $X/M$ .
- (2)  $W + M$  is  $f$ -simultaneously quasi-Chebyshev in  $X$ .

**Proof:**

(1)  $\Rightarrow$  (2) Since  $M$  is  $f$ -simultaneously proximal by Theorem 8,  $W + M$  is  $f$ -simultaneously proximal in  $X$ . Let  $A$  be an arbitrary  $f$ -bounded set in  $X$ . Then  $P_{W+M}^f(A) \neq \phi$ . Now to show that  $P_{W+M}^f(A)$  is  $f$ -compact, we need to show that every sequence in  $P_{W+M}^f(A)$  has an  $f$ -convergent subsequence. Let  $\{g_n\}_{n=1}^\infty$  be an arbitrary sequence in  $P_{W+M}^f(A)$ . Then by Theorem 6, for each  $n > 1$ ,  $g_n + M \in P_{(W+M)/M}^{\tilde{f}}(A/M)$ . Since  $P_{(W+M)/M}^{\tilde{f}}(A/M)$  is  $\tilde{f}$ -compact, one can choose  $g_0 \in W + M$  with  $g_0 + M \in P_{(W+M)/M}^{\tilde{f}}(A/M)$  and  $\{g_{n_k} + M\}_{k=1}^\infty$  is  $\tilde{f}$ -convergent to  $g_0 + M$  for some subsequence  $\{g_{n_k} + M\}_{k=1}^\infty$  of  $\{g_n + M\}_{n=1}^\infty$ . That means,

$$\tilde{f}(g_0 - g_{n_k} + M) = \inf_{m \in M} |f(g_0 - g_{n_k} - m)| \rightarrow 0.$$

Now, since  $M$  is  $f$ -proximal in  $X$ , there exists  $m_{n_k} \in M$  such that  $m_{n_k} \in P_M^f(g_0 - g_{n_k})$ , for every  $k \geq 1$ , and hence

$$|f(g_0 - g_{n_k} - m_{n_k})| = \inf_{m \in M} |f(g_0 - g_{n_k} - m)|.$$

Therefore,

$$\lim_{k \rightarrow \infty} f(g_0 - g_{n_k} - m_{n_k}) = 0.$$

On the other hand,  $\{g_{n_k}\}_{k=1}^\infty$  is an  $f$ -bounded sequence because  $g_n \in P_{W+M}^f(A)$ . In fact  $|f(g_n)| \leq 2 \sup_{a \in A} |f(a)|$ . Since  $M$  has the  $f$ -property, with out loss of generality, we may assume

that for some  $m_0 \in M$ ,  $f(m_{n_k} - m_0) \rightarrow 0$ . Let  $g' = g_0 - m_0$ . Then,  $g' \in W + M$  and

$$\begin{aligned} f(g' - g_{n_k}) &= f(g_0 - m_0 - g_{n_k}) \\ &\leq f(g_0 - g_{n_k} - m_{n_k}) + f(m_{n_k} - m_0), \end{aligned}$$

$\forall k \geq 1$ . Thus,  $\lim_{k \rightarrow \infty} f(g' - g_{n_k}) = 0$ . Since  $\{g_{n_k}\}_{k=1}^{\infty} \in P_{W+M}^f(A)$ , for every  $k \geq 1$ , and  $P_{W+M}^f(A)$  is  $f$ -closed, since  $W + M$  is  $f$ -closed by Proposition 19, we conclude that  $g' \in P_{W+M}^f(A)$ . Hence,  $P_{W+M}^f(A)$  is  $f$ -compact.

(2)  $\Rightarrow$  (1) Since  $M$  and  $W + M$  are  $f$ -simultaneously proximal and  $M \subseteq W + M$ , then  $(W + M)/M = W/M$  is  $\tilde{f}$ -simultaneously proximal in  $X/M$ .

Now, let  $A$  be an arbitrary  $f$ -bounded set in  $X$ . Then,  $P_{W/M}^{\tilde{f}}(A/M)$  is non-empty. So from the hypothesis we have  $W + M$  is  $f$ -simultaneously quasi-Chebyshev in  $X$ , and hence  $P_{W+M}^f(A)$  is  $f$ -compact in  $X$ . Using Theorem 6, we conclude that

$$P_{(W+M)/M}^{\tilde{f}}(A/M) = \pi \left( P_{W+M}^f(A) \right),$$

where  $\pi : X \rightarrow X/M$ ,  $\pi(x) = x + M$ , is continuous. Consequently  $P_{W/M}^{\tilde{f}}(A/M)$  is  $\tilde{f}$ -compact. Therefore,  $W/M$  is  $f$ -simultaneously quasi-Chebyshev in  $X$ .  $\square$

Note that Theorem 11 is still true if the restriction  $W + M$  is  $f$ -closed is replaced by the condition that the function  $f(x) = 0$  if and only if  $x = 0$  and use Theorem 4 to prove that  $W + M$  is  $f$ -closed.

#### 4. Conclusions

In this paper we introduce and study the concept of  $f$ -simultaneous approximation of a nonempty subset  $K$  of Hausdorff topological vector space  $X$ , existence and uniqueness as a generalization to the problem of simultaneous approximation in the sense that if the function  $f$  is taken to be the usual norm, the problem is turned out to be precisely the problem of best approximation in the usual sense. Further, we obtain some results regarding  $f$ -simultaneous approximation in the quotient space.

#### REFERENCES

- Breckner, W. W. and Brosowski, B. (1971). Ein kriterium zur charakterisierung von sonnen, *Mathematika*, Vol. 13, pp. 181-188.
- Govindarajulu, P. and Pai, D. V. (1980). On properties of sets related to  $f$ -projections, *J. Math. Anal. Appl.* Vol. 73, pp. 65-457.
- Khalil, R. and Abushamla, W. (2003).  $f$ -proximality in function spaces, *Dirasat*, Vol. 30, pp. 93-96.
- Moghaddam, M. A. (2010). On  $f$ -best approximation in quotient topological vector spaces, *Int. Math. Forum*. Vol. 12, pp. 587-595.

- Narang, T. D. (1985). On  $f$ -best approximation in topological spaces, Arch. Math. Vol. 4, pp. 229-234.
- Narang, T. D. (1986). Approximation relative to an ultra function, Arch. Math., Vol. 4, pp. 181-186.
- Pai, D. V. and Veermani, P. (1982). Applications of fixed point theorems in optimization and best approximation, Nonlinear Anal. Appl. Marcel Dekker, Int. New York, Edited by S.P. Singh and J. H. Barry, pp. 393-400.
- Rezapour, Sh. (2003).  $\epsilon$ -weakly Chebyshev subspaces of Banach spaces, Anal. Theory Appl., Vol.19, No. 2, pp. 130-135.
- Rezapour, Sh. and Mohebi, H. (2003).  $\epsilon$ -weakly Chebyshev subspaces and quotient spaces, Bulletin of the Iranian Math. Soc., Vol. 29, No. 2, pp. 27-33.
- Saidi, F., Hussein, D. and Khalil, R. (2002). Best simultaneous approximation in  $L^p(I, X)$ , Approx. Theory J., Vol. 116, pp. 369-379.
- Singer, I. (1970). Best approximation in normed linear spaces by elements of linear subspaces, Springer-Verlag.